

Datasheet for ABIN6284789
anti-BAD antibody (AA 70-150)



[Go to Product page](#)

Overview

Quantity:	50 µL
Target:	BAD
Binding Specificity:	AA 70-150
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BAD antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)

Product Details

Purpose:	Bad is a protein encoded by the BAD gene which is approximately 18,4 kDa. Bad is localised to the mitochondrion outer membrane and cytoplasm. It is involved in RET signalling, common cytokine receptor gamma-chain family signalling pathways, glioma and the development HGF signalling pathway. It positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. It also appears to act as a link between growth factor receptor signalling and the apoptotic pathways. Bad is widely expressed in a wide variety of tissues. Mutations in the BAD gene may result in adrenal cortical adenocarcinoma and endometrial cancer. STJ91800 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. This polyclonal antibody detects endogenous levels of Bad protein.
Immunogen:	Synthesized peptide derived from human Bad around the non-phosphorylation site of S134.

Product Details

Isotype:	IgG
Specificity:	Bad Polyclonal Antibody detects endogenous levels of Bad protein.
Characteristics:	Rabbit polyclonal to Bad.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Target Details

Target:	BAD
Alternative Name:	Bad (BAD Products)
Molecular Weight:	28 kDa
Gene ID:	572
UniProt:	Q92934
Pathways:	MAPK Signaling , PI3K-Akt Signaling , RTK Signaling , Apoptosis , Fc-epsilon Receptor Signaling Pathway , Positive Regulation of Peptide Hormone Secretion , Carbohydrate Homeostasis , Positive Regulation of Endopeptidase Activity , Regulation of Carbohydrate Metabolic Process , Hepatitis C , CXCR4-mediated Signaling Events

Application Details

Application Notes:	WB 1:500-1:2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:5000
Comment:	Expressed in a wide variety of tissues.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.
Preservative:	Sodium azide

Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C, and avoid repeat freeze-thaw cycles.