antibodies -online.com





anti-MLX antibody (Internal Region)

MLX

Mlx (MLX Products)



Go to Product page

| \sim | | | |
|--------|-----|------|-----|
| | N/P | r\/I | i⊢₩ |

Target:

Alternative Name:

| Quantity: | 50 μL |
|----------------------|---|
| Target: | MLX |
| Binding Specificity: | AA 80-160, Internal Region |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MLX antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |
| Product Details | |
| Purpose: | Rabbit polyclonal to Mlx. |
| Immunogen: | Synthesized peptide derived from human Mlx |
| Isotype: | IgG |
| Specificity: | Mlx Polyclonal Antibody detects endogenous levels of Mlx protein. |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| | |

Target Details

| Molecular Weight: | 33 kDa | |
|-------------------|--------|--|
| Gene ID: | 6945 | |
| UniProt: | Q9UH92 | |

Application Details

| Application Notes: | WB 1:500-1:2000 |
|--------------------|---|
| | IHC 1:100-1:300 |
| | ELISA 1:10000 |
| Comment: | Expressed in all tissues tested, including spleen, thymus, prostate, ovary, intestine, colon, |
| | peripheral blood leukocyte, heart, liver, skeletal muscle and kidney. Lower levels of expression in |
| | testis, brain, placenta and lung. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C, and avoid repeat freeze-thaw cycles. |