# antibodies -online.com





# anti-RHOH antibody (C-Term)



Go to Product page

| $\sim$ |       |
|--------|-------|
| ( )\/白 | rview |
| OVC    |       |

| Quantity:            | 50 μL   |
|----------------------|---|
| Target:              | RHOH  |
| Binding Specificity: | AA 110-190, C-Term  |
| Reactivity:          | Human, Mouse, Rat   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This RHOH antibody is un-conjugated   |
| Application:         | Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)) |

#### **Product Details**

| Purpose:      | Rabbit polyclonal to Rho H.   |
|---------------|---|
| Immunogen:    | Synthesized peptide derived from human Rho H  |
| Isotype:      | IgG   |
| Specificity:  | Rho H Polyclonal Antibody detects endogenous levels of Rho H protein.   |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

## Target Details

|--|

## Target Details

Storage:

Storage Comment:

| Alternative Name: | Rho H (RHOH Products) |
|-------------------|-----------------------|
| Molecular Weight: | 21 kDa                |
| Gene ID:          | 399                   |
| UniProt:          | Q15669                |

| UniProt:            | Q15669   |
|---------------------|--|
| Application Details |  |
| Application Notes:  | WB 1:500-1:2000  |
|                     | IHC 1:100-1:300  |
|                     | IF 1:200-1:1000  |
|                     | ELISA 1:20000  |
| Comment:            | Expressed only in hematopoietic cells. Present at very high levels in the thymus, less abundan |
|                     | in the spleen, and least abundant in the bone marrow. Expressed at a higher level in the TH1   |
|                     | subtype of T-helper cells than in the TH2 subpopulation. Expressed in neutrophils under        |
|                     | inflammatory conditions, such as cystic fibrosis, ulcerative colitis and appendicitis.         |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Concentration:      | 1 mg/mL  |
| Buffer:             | Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.                     |
| Preservative:       | Sodium azide   |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                  |

should be handled by trained staff only.

Store at -20°C, and avoid repeat freeze-thaw cycles.

-20 °C