# .-online.com antibodies

Datasheet for ABIN6291084 anti-ZNF12 antibody

Image



### Overview

Quantity:	50 µL
Target:	ZNF12
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF12 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

## Product Details

Immunogen:	Recombinant protein of human ZNF12
lsotype:	lgG
Purification:	Affinity purification

## Target Details

Target:	ZNF12
Alternative Name:	ZNF12 (ZNF12 Products)
Background:	This gene is a member of the krueppel C2H2-type zinc-finger protein family and encodes a protein with eight C2H2-type zinc fingers and a KRAB domain. This nuclear protein is involved in developmental control of gene expression. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
Molecular Weight:	81.202 kDa

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN6291084 | 09/10/2023 | Copyright antibodies-online. All rights reserved.


## Target Details

Gene ID:	7559	
UniProt:	P17014	

# Application Details

Application Notes:	WB 1:500 - 1:2000
	IHC 1:50 - 1:200
Comment:	Widely expressed in various adult tissues and embryonic developmental stages (isoform 3)
Restrictions:	For Research Use only
Handling	
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20C. Avoid freeze / thaw cycles.

### Images



### Western Blotting

**Image 1.** Western blot analysis of extracts of mouse kidney, using ZNF12 Antibody (ABIN6291084) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.