

## Datasheet for ABIN6294242 anti-FBXO5 antibody (AA 1-250)



## Overview

0.0	
Quantity:	100 μg
Target:	FBXO5
Binding Specificity:	AA 1-250
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FBXO5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS), Immunohistochemistry (IHC)
Product Details	
Purpose:	Mouse anti-Human EMI1 Antibody [Sodium Azide Free]
Immunogen:	A recombinant fragment (203 amino acid residues between aa 1-250) from the human protein was used as the immunogen for the EMI1 antibody.
Specificity:	Nuclear
Purification:	A recombinant fragment (203 amino acid residues between aa 1-250) from the human protein was used as the immunogen for the EMI1 antibody.
Target Details	
Target:	FBXO5
Alternative Name:	F-box only protein 5 (FBXO5 Products)

## **Target Details**

-	
Background:	Target Description: Early Mitotic Inhibitor-1 regulates mitosis by inhibiting the anaphase
	promoting complex/cyclosome (APC). It is a conserved F box protein containing a zinc-binding
	region essential for APC inhibition. The protein functions to promote cyclin A accumulation and
	S phase entry in somatic cells by inhibiting the APC complex. At the G1-S transition, EMI1 is
	transcriptionally induced by the E2F transcription factor. Overexpression accelerates S-phase
	entry and can override a G1 block caused by overexpression of Cdh1 or the E2F-inhibitor p105
	retinoblastoma protein (pRb). Depleting cells of EMI1 through RNA interference prevents
	accumulation of cyclin A and inhibits S phase entry.
	Gene Symbol: FBXO5///EMI1///FBX5
Gene ID:	26271
UniProt:	Q9UKT4
Pathways:	Mitotic G1-G1/S Phases
Application Details	
Application Notes:	Flow Cytometry: 0.5-1 μg/million cells in 0.1ml
	Immunofluorescence: 0.5-1 μg/mL
	Western blot: 1-2 μg/mL
	Immunohistochemistry (FFPE): 0.5-1 μg/mL for 30 minutes at RT (1)
	Prediluted format: incubate for 30 min at RT (2)
Restrictions:	For Research Use only
Handling	
Buffer:	In 1X PBS, BSA free, sodium azide free
Preservative:	Azide free
Storage:	4 °C,-20 °C
Storage Comment:	2-8°C. The azide-free format should be aliquoted and stored at -20°C or colder.