

Datasheet for ABIN6296037 anti-RNMT antibody



Overview

Quantity:	100 µg
Target:	RNMT
Reactivity:	Human, Rat, Cow, Dog, Hamster
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This RNMT antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunofluorescence (IF), Flow Cytometry (FACS)
Product Details	
Product Details Purpose:	Mouse anti-Human/Cow/Dog/Hamster/Rat TAG-72 Antibody [Sodium Azide Free]
	Mouse anti-Human/Cow/Dog/Hamster/Rat TAG-72 Antibody [Sodium Azide Free] A membrane-enriched fraction of a human breast carcinoma liver metastasis was used as the immunogen for this TAG-72 antibody.
Purpose:	A membrane-enriched fraction of a human breast carcinoma liver metastasis was used as the

Target Details

Target:	RNMT
Alternative Name:	mRNA cap guanine-N7 methyltransferase (RNMT Products)
Background:	Target Description: Recognizes an oncofetal antigen of 220 kDa, identified as a Tumor-
	associated glycoprotein (TAG-72) with properties of a mucin. This mAb defines the mucin-

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN6296037 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

	carried sialylated-Tn epitope. TAG72 is usually expressed by adenocarcinomas, but is negative
	in mesotheliomas. Studies have reported that this antibody has 80 % sensitivity and 93 %
	specificity for pulmonary adenocarcinoma. Therefore, TAG72 is a useful marker to distinguish
	between mesothelioma and adenocarcinoma. However, false positive reactions can occur so
	results must be interpreted with the utmost caution. This antibody may be useful in the
	differentiation of non-small cell carcinomas from small cell carcinomas of the lung. The
	combined use of anti-TAG72 and anti-GCDFP-15 is valuable in the diagnosis of apocrine
	carcinoma.
	Gene Symbol: tag-72 C25A1.3
Gene ID:	182875
UniProt:	Q9XVS1
Application Details	
Application Notes:	Flow Cytometry: 0.5-1 µg/million cells in 0.1ml
Application Notes:	Flow Cytometry: 0.5-1 µg/million cells in 0.1ml Immunofluorescence: 1-2 µg/mL
Application Notes:	
Application Notes:	Immunofluorescence: 1-2 µg/mL
Application Notes: Restrictions:	Immunofluorescence: 1-2 μg/mL Immunohistochemistry (FFPE): 0.5-1 μg/mL for 30 minutes at RT (1)
Restrictions:	Immunofluorescence: 1-2 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format: incubate for 30 min at RT (2)
	Immunofluorescence: 1-2 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format: incubate for 30 min at RT (2)
Restrictions:	Immunofluorescence: 1-2 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format: incubate for 30 min at RT (2)
Restrictions: Handling	Immunofluorescence: 1-2 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format: incubate for 30 min at RT (2) For Research Use only
Restrictions: Handling Buffer:	Immunofluorescence: 1-2 µg/mL Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1) Prediluted format: incubate for 30 min at RT (2) For Research Use only In 1X PBS, BSA free, sodium azide free