

Datasheet for ABIN6296497

anti-CD57 antibody



Go to Product page

_						
	W	0	rv	10	W	

Quantity:	100 μg	
Target:	CD57 (B3GAT1)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CD57 antibody is un-conjugated	
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), Immunofluorescence (IF)	
Product Details		
Purpose:	Mouse anti-Human CD57 Antibody Cocktail [Sodium Azide Free]	
Immunogen:	Human peripheral blood mononuclear cells were used as the immunogen for the CD57 antibody cocktail.	
Specificity:	Cell surface, cytoplasmic	
Purification:	Human peripheral blood mononuclear cells were used as the immunogen for the CD57 antibody cocktail.	
Target Details		
Target:	CD57 (B3GAT1)	
Alternative Name:	Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1 (B3GAT1 Products)	
Background:	Target Description: Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells.	

Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles.

Target Details

rarget Details		
	Anti-CD57 also stains neuroendocrine cells and their derived tumors, including carcinoid tumor and medulloblastoma. Anti-CD57 can also be useful in separating type B3 thymoma from thymic carcinoma when combined with a panel that includes antibodies against GLUT1, CD5, and CEA. Gene Symbol: B3GAT1///GLCATP	
Gene ID:	27087	
UniProt:	Q9P2W7	
Pathways:	Glycosaminoglycan Metabolic Process	
Application Details		
Application Notes:	FACS: 0.5-1 µg/million cells in 0.1ml Immunofluorescence: 0.5-1 µg/mL	
	Immunohistochemistry (FFPE): 2-4 µg/mL for 30 min at RT (1)	
	Prediluted format: incubate for 30 min at RT (2)	
Restrictions:	For Research Use only	
Handling		
Buffer:	In 1X PBS, BSA free, sodium azide free	
Preservative:	Azide free	

4 °C,-20 °C

Storage:

2-8°C. The azide-free format should be aliquoted and stored at -20°C or colder.