

Datasheet for ABIN630031
anti-HNRPLL antibody (N-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	HNRPLL
Binding Specificity:	N-Term
Reactivity:	Human, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRPLL antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	HNRPLL antibody was raised using the N terminal of HNRPLL corresponding to a region with amino acids RLKTEEGEIDYSAEEGENRREATPRGGGDGGGGGRSFSQPEAGGSHHKVS
Specificity:	HNRPLL antibody was raised against the N terminal of HNRPLL
Purification:	Purified

Target Details

Target:	HNRPLL
Alternative Name:	HNRPLL (HNRPLL Products)
Background:	HNRPLL contains 3 RRM (RNA recognition motif) domains and may bind RNA and plays a role in mRNA processing.

Target Details

Molecular Weight: 60 kDa (MW of target protein)

Application Details

Application Notes: WB: 1.25 µg/mL, IHC: 4-8 µg/mL
Optimal conditions should be determined by the investigator.

Comment: HNRPLL Blocking Peptide, catalog no. 33R-8037, is also available for use as a blocking control in assays to test for specificity of this HNRPLL antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Lyophilized powder. Add distilled water for a 1 mg/mL concentration of HNRPLL antibody in PBS

Concentration: Lot specific


Buffer: PBS

Handling Advice: Avoid repeated freeze/thaw cycles.
Dilute only prior to immediate use.

Storage: 4 °C/-20 °C


Storage Comment: Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images




Immunohistochemistry

Image 1. HNRPLL antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Alveolar cells (arrows) in Human Lung. Magnification is at 400X.



Western Blotting

Image 2. HNRPLL antibody used at 1.25 ug/ml to detect target protein.



Immunohistochemistry

Image 3. HNRPLL antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Myocardial cells (arrows) in Human Heart. Magnification is at 400X