

Datasheet for ABIN630077 anti-KCTD18 antibody (N-Term)

1 Image



Go to Product page

/
١

Quantity:	100 μg
Target:	KCTD18
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCTD18 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	KCTD18 antibody was raised using the N terminal of KCTD18 corresponding to a region with
	KCTD18 antibody was raised using the N terminal of KCTD18 corresponding to a region with amino acids LHYLNTSGASCESRIIGVYATKTDGTDAIEKQLGGRIHSKGIFKREAGNN
Immunogen:	amino acids LHYLNTSGASCESRIIGVYATKTDGTDAIEKQLGGRIHSKGIFKREAGNN
Immunogen: Specificity:	amino acids LHYLNTSGASCESRIIGVYATKTDGTDAIEKQLGGRIHSKGIFKREAGNN KCTD18 antibody was raised against the N terminal of KCTD18
Immunogen: Specificity: Purification:	amino acids LHYLNTSGASCESRIIGVYATKTDGTDAIEKQLGGRIHSKGIFKREAGNN KCTD18 antibody was raised against the N terminal of KCTD18
Immunogen: Specificity: Purification: Target Details	amino acids LHYLNTSGASCESRIIGVYATKTDGTDAIEKQLGGRIHSKGIFKREAGNN KCTD18 antibody was raised against the N terminal of KCTD18 Purified
Immunogen: Specificity: Purification: Target Details Target:	amino acids LHYLNTSGASCESRIIGVYATKTDGTDAIEKQLGGRIHSKGIFKREAGNN KCTD18 antibody was raised against the N terminal of KCTD18 Purified KCTD18

Application Details

Application Notes:	WB: 5 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	KCTD18 Blocking Peptide, catalog no. 33R-5035, is also available for use as a blocking control in assays to test for specificity of this KCTD18 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of KCTD18 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. KCTD18 antibody used at 5 ug/ml to detect target protein.