

Datasheet for ABIN630368
anti-SLC35F2 antibody (N-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	SLC35F2
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC35F2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	SLC35 F2 antibody was raised using the N terminal of SLC35 2 corresponding to a region with amino acids MEADSPAGPGAPEPLAEGAAAEFSSLLRRRIKGLFTWNILKTIALGQMLS
Specificity:	SLC35 F2 antibody was raised against the N terminal of SLC35 2
Purification:	Purified

Target Details

Target:	SLC35F2
Alternative Name:	SLC35F2 (SLC35F2 Products)
Background:	SLC35F2 is a putative solute transporter.
Molecular Weight:	41 kDa (MW of target protein)


Application Details

Application Notes:	WB: 2.5 µg/mL, IHC: 4-8 µg/mL Optimal conditions should be determined by the investigator.
Comment:	SLC35F2 Blocking Peptide, catalog no. 33R-5885, is also available for use as a blocking control in assays to test for specificity of this SLC35F2 antibody
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of SLC30 2 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images




Immunohistochemistry

Image 1. SLC35F2 antibody was used for immunohistochemistry at a concentration of 4-8 µg/ml. Magnification is at 400X



Western Blotting

Image 2. SLC35F2 antibody used at 2.5 ug/ml to detect target protein.



Immunohistochemistry

Image 3. SLC35F2 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Skeletal muscle cells (arrows) in Human Muscle. Magnification is at 400X