

Datasheet for ABIN630475 anti-MBOAT7 antibody (C-Term)

1 Image



Go to Product page

| \sim | | | | |
|--------|----------------|-------|--------|----|
| () | Ive | r\ / | \cap | Λ. |
| \cup | $\lor \subset$ | I V I | \Box | ٧V |

| Quantity: | 100 μg |
|----------------------|---|
| Target: | MBOAT7 |
| Binding Specificity: | C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MBOAT7 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | LENG4 antibody was raised using the C terminal Of Leng4 corresponding to a region with |
| | amino acids LSAYWHGLHPGYYLSFLTIPLCLAAEGRLESALRGRLSPGGQKAWDWVHW |
| Specificity: | LENG4 antibody was raised against the C terminal Of Leng4 |
| Purification: | Purified |
| Target Details | |
| Target: | MBOAT7 |
| Alternative Name: | LENG4 (MBOAT7 Products) |
| Background: | The function of LENG4 protein is not widely studied, and is yet to be elucidated fully. |
| Molecular Weight: | 53 kDa (MW of target protein) |
| | |

Target Details

| Pathways: Ir | nositol Metabolic Process |
|--------------|---------------------------|
|--------------|---------------------------|


Application Details

| Application Notes: | WB: 5 µg/mL |
|--------------------|--|
| | Optimal conditions should be determined by the investigator. |
| Comment: | LENG4 Blocking Peptide, catalog no. 33R-5412, is also available for use as a blocking control in assays to test for specificity of this LENG4 antibody |
| Restrictions: | For Research Use only |

Handling

| Format: | Lyophilized | |
|------------------|--|--|
| Reconstitution: | Lyophilized powder. Add distilled water for a 1 mg/mL concentration of LENG4 antibody in PBS | |
| Concentration: | Lot specific | |
| Buffer: | PBS | |
| Handling Advice: | Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C. | |

Images



Western Blotting

Image 1. LENG4 antibody used at 5 ug/ml to detect target protein.