

## Datasheet for ABIN630508 **anti-NR2F1 antibody (N-Term)**



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	100 µL
Target:	NR2F1
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR2F1 antibody is un-conjugated
Application:	Western Blotting (WB)

#### Product Details

Immunogen:	NR2 F1 antibody was raised using the N terminal of NR2 1 corresponding to a region with amino acids GEQQQAGSGAPHTPQTPGQPGAPATPGTAGDKGQGGPPGSGQSQQHIECV
Specificity:	NR2 F1 antibody was raised against the N terminal of NR2 1
Purification:	Affinity purified

#### Target Details

Target:	NR2F1
Alternative Name:	NR2F1 ( <a href="#">NR2F1 Products</a> )
Background:	Coup (chicken ovalbumin upstream promoter) transcription factor binds to the ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. NR2F1 binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif.

## Target Details

Molecular Weight: 46 kDa (MW of target protein)

Pathways: Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway

## Application Details

Application Notes: WB: 1 µg/mL

Optimal conditions should be determined by the investigator.

Comment: NR2F1 Blocking Peptide, catalog no. 33R-3245, is also available for use as a blocking control in assays to test for specificity of this NR2F1 antibody

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Reconstitution: Lyophilized powder. Add distilled water for a 1 mg/mL concentration of NR0 1 antibody in PBS

Concentration: Lot specific

Buffer: PBS

Handling Advice: Avoid repeated freeze/thaw cycles.  
Dilute only prior to immediate use.

Storage: 4 °C/-20 °C

Storage Comment: Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

## Images



### Western Blotting

**Image 1.** NR2F1 antibody used at 1 µg/ml to detect target protein.