

Datasheet for ABIN630695 anti-GAPVD1 antibody (N-Term)





\sim			
	ve	r\/	٨
\cup	V C	1 V I	٧V

100 μL	
GAPVD1	
Human, Mouse, Rat, Dog	
Rabbit	
Polyclonal	
This GAPVD1 antibody is un-conjugated	
Western Blotting (WB), Immunohistochemistry (IHC)	
esponding to a region with	
KLIERFSPSQQ	
anine nucleotide exchange	
anine nucleotide exchange vtosis, insulin receptor	

by exchanging bound GDP for free GTP, leading to regulate LC2A4/GLUT4 trafficking. In the absence of insulin, it maintains RAB31 in an active state and promotes a futile cycle between LC2A4/GLUT4 storage vesicles and early endosomes, retaining LC2A4/GLUT4 inside the cells. Upon insulin stimulation, it is translocated to the plasma membrane, releasing LC2A4/GLUT4 from intracellular storage vesicles. It is also involved in EGFR trafficking and degradation, possibly by promoting EGFR ubiquitination and subsequent degradation by the proteasome. It has GEF activity for Rab5 and GAP activity for Ras.

Molecular Weight:


166 kDa (MW of target protein)

Application Details

Application Notes:	WB: 0.25 μg/mL, IHC: 4-8 μg/mL		
	Optimal conditions should be determined by the investigator.		
Comment:	GAPVD1 Blocking Peptide, catalog no. 33R-2950, is also available for use as a blocking control in assays to test for specificity of this GAPVD1 antibody		
Restrictions:	For Research Use only		


Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of GAPVD1 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	




Immunohistochemistry

Image 1. GAPVD1 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml. Magnification is at 400X



Western Blotting

Image 2. GAPVD1 antibody used at 0.25 ug/ml to detect target protein.



Immunohistochemistry

Image 3. GAPVD1 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Neural cells (arrows) in Human Brain. Magnification is at 400X