

Datasheet for ABIN631192 anti-NGRN antibody (N-Term)

1 Image



Go to Product page

()	ve	V /	-	1 A
	\cup	1 \/	-	1/1
\sim	' V C	1 V	ı	v v

Quantity:	100 μL
Target:	NGRN
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NGRN antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	NGRN antibody was raised using the N terminal of NGRN corresponding to a region with amino
	NGRN antibody was raised using the N terminal of NGRN corresponding to a region with amino acids MAVTLSLLLGGRVCAAVTRCGFATRGVAGPGPIGREPDPDSDWEPEEREL
Immunogen:	acids MAVTLSLLLGGRVCAAVTRCGFATRGVAGPGPIGREPDPDSDWEPEEREL
Immunogen: Specificity:	acids MAVTLSLLLGGRVCAAVTRCGFATRGVAGPGPIGREPDPDSDWEPEEREL NGRN antibody was raised against the N terminal of NGRN
Immunogen: Specificity: Purification:	acids MAVTLSLLLGGRVCAAVTRCGFATRGVAGPGPIGREPDPDSDWEPEEREL NGRN antibody was raised against the N terminal of NGRN
Immunogen: Specificity: Purification: Target Details	acids MAVTLSLLLGGRVCAAVTRCGFATRGVAGPGPIGREPDPDSDWEPEEREL NGRN antibody was raised against the N terminal of NGRN Affinity purified
Immunogen: Specificity: Purification: Target Details Target:	acids MAVTLSLLLGGRVCAAVTRCGFATRGVAGPGPIGREPDPDSDWEPEEREL NGRN antibody was raised against the N terminal of NGRN Affinity purified NGRN


Application Details

Application Notes:	n Notes: WB: 1 μg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	NGRN Blocking Peptide, catalog no. 33R-5804, is also available for use as a blocking control in assays to test for specificity of this NGRN antibody	
Restrictions:	For Research Use only	

Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of NGRN antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

Images



Western Blotting

Image 1. NGRN antibody used at 1 ug/ml to detect target protein.