

Datasheet for ABIN631248 **anti-WASF3 antibody (N-Term)**



[Go to Product page](#)

1 Image

Overview

| | |
|----------------------|--------------------------------------|
| Quantity: | 100 µL |
| Target: | WASF3 |
| Binding Specificity: | N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This WASF3 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|---------------|--|
| Immunogen: | WASF3 antibody was raised using the N terminal of WASF3 corresponding to a region with amino acids NMKKAFKSSTVQDQQVSKNSIPNPVADIYNQSDKPPPLNILTPYRDDKK |
| Specificity: | WASF3 antibody was raised against the N terminal of WASF3 |
| Purification: | Affinity purified |

Target Details

| | |
|-------------------|---|
| Target: | WASF3 |
| Alternative Name: | WASF3 (WASF3 Products) |
| Background: | WASF3 is a member of the Wiskott-Aldrich syndrome protein family. It is a protein that forms a multiprotein complex that links receptor kinases and actin. Binding to actin occurs through a C-terminal verprolin homology domain in all family members. The multiprotein complex serves to |

Target Details

transduce signals that involve changes in cell shape, motility or function.

Molecular Weight: 55 kDa (MW of target protein)

Pathways: [RTK Signaling](#)

Application Details

Application Notes: WB: 1 µg/mL

Optimal conditions should be determined by the investigator.

Comment: WASF3 Blocking Peptide, catalog no. 33R-6796, is also available for use as a blocking control in assays to test for specificity of this WASF3 antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Lyophilized powder. Add distilled water for a 1 mg/mL concentration of WASF3 antibody in PBS

Concentration: Lot specific

Buffer: PBS

Handling Advice: Avoid repeated freeze/thaw cycles.
Dilute only prior to immediate use.

Storage: 4 °C/-20 °C

Storage Comment: Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.



Western Blotting

Image 1. WASF3 antibody used at 1 ug/ml to detect target protein.