

# Datasheet for ABIN631543

# anti-TTC16 antibody (C-Term)





Go to Product page

0				

Quantity:	100 μL
Target:	TTC16
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TTC16 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details  Immunogen:	TTC16 antibody was raised using the C terminal of TTC16 corresponding to a region with
	TTC16 antibody was raised using the C terminal of TTC16 corresponding to a region with amino acids RSRGLLRSSTKTEAFYDSNWSLSKTEYAQGQGQRSSKAEGAQGKSQGMSS
Immunogen:	amino acids RSRGLLRSSTKTEAFYDSNWSLSKTEYAQGQGQRSSKAEGAQGKSQGMSS
Immunogen: Specificity:	amino acids RSRGLLRSSTKTEAFYDSNWSLSKTEYAQGQGQRSSKAEGAQGKSQGMSS  TTC16 antibody was raised against the C terminal of TTC16
Immunogen:  Specificity:  Purification:	amino acids RSRGLLRSSTKTEAFYDSNWSLSKTEYAQGQGQRSSKAEGAQGKSQGMSS  TTC16 antibody was raised against the C terminal of TTC16
Immunogen:  Specificity:  Purification:  Target Details	amino acids RSRGLLRSSTKTEAFYDSNWSLSKTEYAQGQGQRSSKAEGAQGKSQGMSS  TTC16 antibody was raised against the C terminal of TTC16  Affinity purified
Immunogen:  Specificity:  Purification:  Target Details  Target:	amino acids RSRGLLRSSTKTEAFYDSNWSLSKTEYAQGQGQRSSKAEGAQGKSQGMSS  TTC16 antibody was raised against the C terminal of TTC16  Affinity purified  TTC16


# **Application Details**

Application Notes:	WB: 1 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	TTC16 Blocking Peptide, catalog no. 33R-8209, is also available for use as a blocking control in
	assays to test for specificity of this TTC16 antibody
Restrictions:	For Research Use only

# Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of TTC16 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles.  Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

### **Images**



### **Western Blotting**

**Image 1.** TTC16 antibody used at 1 ug/ml to detect target protein.