

Datasheet for ABIN631726

anti-FBXO28 antibody (Middle Region)





Overview

Overview	
Quantity:	100 μL
Target:	FBXO28
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXO28 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	FBXO28 antibody was raised using the middle region of FBXO28 corresponding to a region with
	amino acids ELERKLREVMESAVGNSSGSGQNEESPRKRKKATEAIDSLRKSKRLRNRK
Specificity:	FBXO28 antibody was raised against the middle region of FBXO28
Purification:	Affinity purified
Target Details	
Target:	FBXO28
Alternative Name:	FBXO28 (FBXO28 Products)
Background:	Members of the F-box protein family, such as FBXO28, are characterized by an approximately
	40-amino acid F-box motif. SCF complexes, formed by SKP1, cullin, and F-box proteins, act as
	protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact

Target Details

with ubiquitination targets through other protein interaction domains. Members of the F-box		
protein family, such as FBX	(028, are characterized by an approximately 40-amino acid F-box	
motif. SCF complexes, form	ned by SKP1, cullin, and F-box proteins, act as protein-ubiquitin	
ligases. F-box proteins interact with SKP1 through the F box, and they interact with		
ubiquitination targets throu	igh other protein interaction domains.	

Molecular Weight:


41 kDa (MW of target protein)

Application Details

Application Notes:	WB: 1 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	FBXO28 Blocking Peptide, catalog no. 33R-2539, is also available for use as a blocking control in assays to test for specificity of this FBXO28 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of FBXO28 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.



Western Blotting

Image 1. FBXO28 antibody used at 1 ug/ml to detect target protein.