

# Datasheet for ABIN631786 anti-DCAF12 antibody (N-Term)

# 1 Image



Go to Product page

$\sim$					
	ve	r\/		٨	
$\cup$	V C	1 V I		٧V	

Quantity:	100 μL
Target:	DCAF12
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DCAF12 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	WDR40 A antibody was raised using the N terminal of WDR40 corresponding to a region with
	WDR40 A antibody was raised using the N terminal of WDR40 corresponding to a region with amino acids ARKVVSRKRKAPASPGAGSDAQGPQFGWDHSLHKRKRLPPVKRSLVYYLK
Immunogen:	amino acids ARKVVSRKRKAPASPGAGSDAQGPQFGWDHSLHKRKRLPPVKRSLVYYLK
Immunogen: Specificity:	amino acids ARKVVSRKRKAPASPGAGSDAQGPQFGWDHSLHKRKRLPPVKRSLVYYLK  WDR40 A antibody was raised against the N terminal of WDR40
Immunogen:  Specificity:  Purification:	amino acids ARKVVSRKRKAPASPGAGSDAQGPQFGWDHSLHKRKRLPPVKRSLVYYLK  WDR40 A antibody was raised against the N terminal of WDR40
Immunogen:  Specificity:  Purification:  Target Details	amino acids ARKVVSRKRKAPASPGAGSDAQGPQFGWDHSLHKRKRLPPVKRSLVYYLK  WDR40 A antibody was raised against the N terminal of WDR40  Affinity purified
Immunogen:  Specificity:  Purification:  Target Details  Target:	amino acids ARKVVSRKRKAPASPGAGSDAQGPQFGWDHSLHKRKRLPPVKRSLVYYLK  WDR40 A antibody was raised against the N terminal of WDR40  Affinity purified  DCAF12


## **Application Details**

Application Notes:	WB: 1 µg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	WDR40A Blocking Peptide, catalog no. 33R-1481, is also available for use as a blocking control	
	in assays to test for specificity of this WDR40A antibody	
Restrictions:	For Research Use only	

## Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of WDR40 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles.  Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

#### **Images**



#### **Western Blotting**

**Image 1.** WDR40A antibody used at 1 ug/ml to detect target protein.