

Datasheet for ABIN632078 anti-TPRG1 antibody (N-Term)

1 Image



Overview

Quantity:	100 μL
Target:	TPRG1
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TPRG1 antibody is un-conjugated
Application:	Western Blotting (WB)
Draduct Dataila	
Product Details	
Immunogen:	FAM79 B antibody was raised using the N terminal Of Fam79 corresponding to a region with
	FAM79 B antibody was raised using the N terminal Of Fam79 corresponding to a region with amino acids DPMPRQISRQSSVTESTLYPNPYHQPYISRKYFATRPGAIETAMEDLKGH
Immunogen:	amino acids DPMPRQISRQSSVTESTLYPNPYHQPYISRKYFATRPGAIETAMEDLKGH
Immunogen: Specificity:	amino acids DPMPRQISRQSSVTESTLYPNPYHQPYISRKYFATRPGAIETAMEDLKGH FAM79 B antibody was raised against the N terminal Of Fam79
Immunogen: Specificity: Purification:	amino acids DPMPRQISRQSSVTESTLYPNPYHQPYISRKYFATRPGAIETAMEDLKGH FAM79 B antibody was raised against the N terminal Of Fam79
Immunogen: Specificity: Purification: Target Details	amino acids DPMPRQISRQSSVTESTLYPNPYHQPYISRKYFATRPGAIETAMEDLKGH FAM79 B antibody was raised against the N terminal Of Fam79 Affinity purified
Immunogen: Specificity: Purification: Target Details Target:	amino acids DPMPRQISRQSSVTESTLYPNPYHQPYISRKYFATRPGAIETAMEDLKGH FAM79 B antibody was raised against the N terminal Of Fam79 Affinity purified TPRG1


Application Details

Application Notes:	WB: 1 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	FAM79B Blocking Peptide, catalog no. 33R-2108, is also available for use as a blocking control
	in assays to test for specificity of this FAM79B antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of FAM70 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. FAM79B antibody used at 1 ug/ml to detect target protein.