

Datasheet for ABIN632151

anti-KRTAP8-1 antibody (Middle Region)





Go to Product page

_				
()	ve.	rv/	101	Λ

O V CI V I C V V		
Quantity:	100 μL	
Target:	KRTAP8-1	
Binding Specificity:	Middle Region	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KRTAP8-1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	KAP8.1 antibody was raised using the middle region of KRTAP8-1 corresponding to a region	
	with amino acids LCDNFPGAVFPGCYWGSYGYPLGYSVGCGYGSTYSPVGYGFGYGYNGCGA	
Specificity:	KAP8.1 antibody was raised against the middle region of KRTAP8-1	
Purification:	Affinity purified	
Target Details		
Target:	KRTAP8-1	
Alternative Name:	KAP8.1 (KRTAP8-1 Products)	
Background:	KRTAP8-1 belongs to the KRTAP type 8 family. In the hair cortex, hair keratin intermediate	
	filaments are embedded in an interfilamentous matrix, consisting of hair keratin-associated	
	proteins (KRTAP), which are essential for the formation of a rigid and resistant hair shaft	

Target Details

ranger Betane		
	through their extensive disulfide bond cross-linking with abundant cysteine residues of hair keratins. The matrix proteins include the high-sulfur and high-glycine-tyrosine keratins.	
Molecular Weight:	7 kDa (MW of target protein)	
Application Details		
Application Notes:	WB: 1 μg/mL Optimal conditions should be determined by the investigator.	
Comment:	KAP8.1 Blocking Peptide, catalog no. 33R-4817, is also available for use as a blocking control in assays to test for specificity of this KAP8.1 antibody	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of KRTAP8-1 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	



Western Blotting

Image 1. KAP8.1 antibody used at 1 ug/ml to detect target protein.