

Datasheet for ABIN632208
anti-HAGH antibody (C-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	HAGH
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HAGH antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	HAGH antibody was raised using the C terminal of HAGH corresponding to a region with amino acids FARHVEPGNAAIREKLAWAKEKYSIGEPTVPSTLAEFTYNPFMRVREKT
Specificity:	HAGH antibody was raised against the C terminal of HAGH
Purification:	Affinity purified

Target Details

Target:	HAGH
Alternative Name:	HAGH (HAGH Products)
Background:	HAGH is classified as a thiolesterase and is responsible for the hydrolysis of S-lactoyl-glutathione to reduced glutathione and D-lactate. The enzyme encoded by this gene is classified as a thiolesterase and is responsible for the hydrolysis of S-lactoyl-glutathione to

Target Details

reduced glutathione and D-lactate.

Molecular Weight: 29 kDa (MW of target protein)

Application Details

Application Notes: WB: 1 µg/mL

Optimal conditions should be determined by the investigator.

Comment: HAGH Blocking Peptide, catalog no. 33R-2847, is also available for use as a blocking control in assays to test for specificity of this HAGH antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Lyophilized powder. Add distilled water for a 1 mg/mL concentration of HAGH antibody in PBS

Concentration: Lot specific


Buffer: PBS

Handling Advice: Avoid repeated freeze/thaw cycles.
Dilute only prior to immediate use.

Storage: 4 °C/-20 °C

Storage Comment: Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. HAGH antibody used at 1 µg/ml to detect target protein.