

# Datasheet for ABIN632250 anti-WDR66 antibody (N-Term)

# 1 Image



#### Overview

Quantity:	100 μL
Target:	WDR66
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDR66 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	WDR66 antibody was raised using the N terminal of WDR66 corresponding to a region with
	WDR66 antibody was raised using the N terminal of WDR66 corresponding to a region with amino acids GELEEKTDRMPQDELGQERRDLEPENREEGQERRVSDIQSKAGISRESLV
Immunogen:	amino acids GELEEKTDRMPQDELGQERRDLEPENREEGQERRVSDIQSKAGISRESLV
Immunogen: Specificity:	amino acids GELEEKTDRMPQDELGQERRDLEPENREEGQERRVSDIQSKAGISRESLV  WDR66 antibody was raised against the N terminal of WDR66
Immunogen:  Specificity:  Purification:	amino acids GELEEKTDRMPQDELGQERRDLEPENREEGQERRVSDIQSKAGISRESLV  WDR66 antibody was raised against the N terminal of WDR66
Immunogen:  Specificity:  Purification:  Target Details	amino acids GELEEKTDRMPQDELGQERRDLEPENREEGQERRVSDIQSKAGISRESLV  WDR66 antibody was raised against the N terminal of WDR66  Affinity purified
Immunogen:  Specificity:  Purification:  Target Details  Target:	amino acids GELEEKTDRMPQDELGQERRDLEPENREEGQERRVSDIQSKAGISRESLV  WDR66 antibody was raised against the N terminal of WDR66  Affinity purified  WDR66


## **Application Details**

Application Notes:	WB: 1 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	WDR66 Blocking Peptide, catalog no. 33R-3234, is also available for use as a blocking control in
	assays to test for specificity of this WDR66 antibody
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of WDR66 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles.  Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

#### **Images**



#### **Western Blotting**

**Image 1.** WDR66 antibody used at 1 ug/ml to detect target protein.