

Datasheet for ABIN632327

anti-TTC33 antibody (N-Term)





Go to Product page

(۱۱/		٢V	Ĺ		۱٨	١.
	, v	\cup	V	1	$\overline{}$	٧	V

Quantity:	100 μL
Target:	TTC33
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TTC33 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	TTC22 entitled was raised using the Nterminal of TTC22 corresponding to a region with
iriiriariogen.	TTC33 antibody was raised using the N terminal of TTC33 corresponding to a region with
minunogen.	amino acids DEGNWLHAIKRRKEILLEGCAEKSKQLKDEGASLAENKRYREAIQKWDEA
Specificity:	
	amino acids DEGNWLHAIKRRKEILLEGCAEKSKQLKDEGASLAENKRYREAIQKWDEA
Specificity:	amino acids DEGNWLHAIKRRKEILLEGCAEKSKQLKDEGASLAENKRYREAIQKWDEA TTC33 antibody was raised against the N terminal of TTC33
Specificity: Purification:	amino acids DEGNWLHAIKRRKEILLEGCAEKSKQLKDEGASLAENKRYREAIQKWDEA TTC33 antibody was raised against the N terminal of TTC33
Specificity: Purification: Target Details	amino acids DEGNWLHAIKRRKEILLEGCAEKSKQLKDEGASLAENKRYREAIQKWDEA TTC33 antibody was raised against the N terminal of TTC33 Affinity purified
Specificity: Purification: Target Details Target:	amino acids DEGNWLHAIKRRKEILLEGCAEKSKQLKDEGASLAENKRYREAIQKWDEA TTC33 antibody was raised against the N terminal of TTC33 Affinity purified TTC33


Application Details

Application Notes:	WB: 1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	TTC33 Blocking Peptide, catalog no. 33R-1903, is also available for use as a blocking control in
	assays to test for specificity of this TTC33 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of TTC33 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. TTC33 antibody used at 1 ug/ml to detect target protein.