

Datasheet for ABIN632550

anti-GDAP2 antibody (N-Term)





Go to Product page

_			
	Ve.	rv	iew

Quantity:	100 μL
Target:	GDAP2
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GDAP2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	GDAP2 antibody was raised using the N terminal of GDAP2 corresponding to a region with
	GDAP2 antibody was raised using the N terminal of GDAP2 corresponding to a region with amino acids CRTGEAKLTKGFNLAARFIIHTVGPKYKSRYRTAAESSLYSCYRNVLQLA
Immunogen:	amino acids CRTGEAKLTKGFNLAARFIIHTVGPKYKSRYRTAAESSLYSCYRNVLQLA
Immunogen: Specificity:	amino acids CRTGEAKLTKGFNLAARFIIHTVGPKYKSRYRTAAESSLYSCYRNVLQLA GDAP2 antibody was raised against the N terminal of GDAP2
Immunogen: Specificity: Purification:	amino acids CRTGEAKLTKGFNLAARFIIHTVGPKYKSRYRTAAESSLYSCYRNVLQLA GDAP2 antibody was raised against the N terminal of GDAP2
Immunogen: Specificity: Purification: Target Details	amino acids CRTGEAKLTKGFNLAARFIIHTVGPKYKSRYRTAAESSLYSCYRNVLQLA GDAP2 antibody was raised against the N terminal of GDAP2 Affinity purified
Immunogen: Specificity: Purification: Target Details Target:	amino acids CRTGEAKLTKGFNLAARFIIHTVGPKYKSRYRTAAESSLYSCYRNVLQLA GDAP2 antibody was raised against the N terminal of GDAP2 Affinity purified GDAP2


Application Details

Application Notes:	WB: 0.0625 μg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	GDAP2 Blocking Peptide, catalog no. 33R-1801, is also available for use as a blocking control in assays to test for specificity of this GDAP2 antibody	
Restrictions:	For Research Use only	

Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of GDAP2 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

Images



Western Blotting

Image 1. GDAP2 antibody used at 0.0625 ug/ml to detect target protein.