

Datasheet for ABIN632728

anti-WDR63 antibody (Middle Region)





Go to Product page

()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Quantity:	100 μL
Target:	WDR63
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDR63 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	WDR63 antibody was raised using the middle region of WDR63 corresponding to a region with
	WDR63 antibody was raised using the middle region of WDR63 corresponding to a region with amino acids EIALQQNEIMNTFIDDWKYLAEEEGTFGDKTDTHLKEYQSFTDLHSPTEK
Immunogen:	amino acids EIALQQNEIMNTFIDDWKYLAEEEGTFGDKTDTHLKEYQSFTDLHSPTEK
Immunogen: Specificity:	amino acids EIALQQNEIMNTFIDDWKYLAEEEGTFGDKTDTHLKEYQSFTDLHSPTEK WDR63 antibody was raised against the middle region of WDR63
Immunogen: Specificity: Purification:	amino acids EIALQQNEIMNTFIDDWKYLAEEEGTFGDKTDTHLKEYQSFTDLHSPTEK WDR63 antibody was raised against the middle region of WDR63
Immunogen: Specificity: Purification: Target Details	amino acids EIALQQNEIMNTFIDDWKYLAEEEGTFGDKTDTHLKEYQSFTDLHSPTEK WDR63 antibody was raised against the middle region of WDR63 Affinity purified
Immunogen: Specificity: Purification: Target Details Target:	amino acids EIALQQNEIMNTFIDDWKYLAEEEGTFGDKTDTHLKEYQSFTDLHSPTEK WDR63 antibody was raised against the middle region of WDR63 Affinity purified WDR63


Application Details

Application Notes:	WB: 1 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	WDR63 Blocking Peptide, catalog no. 33R-2463, is also available for use as a blocking control in
	assays to test for specificity of this WDR63 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of WDR63 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

Images



Western Blotting

Image 1. WDR63 antibody used at 1 ug/ml to detect target protein.