

Datasheet for ABIN633095

anti-CCDC76 antibody (N-Term)





Go to Product page


| () | 11 / | er | ~\ / | | 1 A |
|--------|-------|----------|------|--------|-----|
| | 1 / / | -1 | \/ | - | 1/1 |
| \sim | v | \sim 1 | V | \sim | V V |
| | | | | | |

| Quantity: | 100 μL |
|--|--|
| Target: | CCDC76 |
| Binding Specificity: | N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CCDC76 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | CCDC76 antibody was raised using the N terminal of CCDC76 corresponding to a region with |
| | amino acids QLAKHLKKCNSREKPKPDFYIQDINAGLRDETEIPEQLVPISSLSEEQLE |
| | |
| Specificity: | CCDC76 antibody was raised against the N terminal of CCDC76 |
| Specificity: Purification: | CCDC76 antibody was raised against the N terminal of CCDC76 Affinity purified |
| | |
| Purification: | |
| Purification: Target Details | Affinity purified |
| Purification: Target Details Target: | Affinity purified CCDC76 |

Target Details

| Molecular Weight: | 54 kDa (MW of target protein) | |
|---------------------|--|--|
| Application Details | | |
| Application Notes: | WB: 1 μg/mL | |
| | Optimal conditions should be determined by the investigator. | |
| Comment: | CCDC76 Blocking Peptide, catalog no. 33R-7618, is also available for use as a blocking control | |
| | in assays to test for specificity of this CCDC76 antibody | |
| Restrictions: | For Research Use only | |
| | | |
| Handling | | |
| Format: | Lyophilized | |
| Reconstitution: | Lyophilized powder. Add distilled water for a 1 mg/mL concentration of CCDC76 antibody in | |
| | PBS | |
| Concentration: | Lot specific | |
| Buffer: | PBS | |
| Handling Advice: | Avoid repeated freeze/thaw cycles. | |
| | Dilute only prior to immediate use. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C. | |

Images



Western Blotting

Image 1. CCDC76 antibody used at 1 ug/ml to detect target protein.