

Datasheet for ABIN633116 anti-RFTN2 antibody (N-Term)

Image



\sim	
()ver	view
0 , 0	* 1 0 * *

100 μL
RFTN2
N-Term
Human
Rabbit
Polyclonal
This RFTN2 antibody is un-conjugated
Western Blotting (WB)
RFTN2 antibody was raised using the N terminal of RFTN2 corresponding to a region with
RFTN2 antibody was raised using the N terminal of RFTN2 corresponding to a region with amino acids MGCGLRKLEDPDDSSPGKIFSTLKRPQVETKTEFAYEYVLLDFTLQASSN
amino acids MGCGLRKLEDPDDSSPGKIFSTLKRPQVETKTEFAYEYVLLDFTLQASSN
amino acids MGCGLRKLEDPDDSSPGKIFSTLKRPQVETKTEFAYEYVLLDFTLQASSN RFTN2 antibody was raised against the N terminal of RFTN2
amino acids MGCGLRKLEDPDDSSPGKIFSTLKRPQVETKTEFAYEYVLLDFTLQASSN RFTN2 antibody was raised against the N terminal of RFTN2
amino acids MGCGLRKLEDPDDSSPGKIFSTLKRPQVETKTEFAYEYVLLDFTLQASSN RFTN2 antibody was raised against the N terminal of RFTN2 Affinity purified
amino acids MGCGLRKLEDPDDSSPGKIFSTLKRPQVETKTEFAYEYVLLDFTLQASSN RFTN2 antibody was raised against the N terminal of RFTN2 Affinity purified RFTN2


Application Details

Application Notes:	WB: 1 µg/mL
	Optimal conditions should be determined by the investigator.
Comment:	RFTN2 Blocking Peptide, catalog no. 33R-6014, is also available for use as a blocking control in assays to test for specificity of this RFTN2 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of RFTN2 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. RFTN2 antibody used at 1 ug/ml to detect target protein.