

Datasheet for ABIN633124 anti-TMEM139 antibody (N-Term)

1 Image



Go to Product page

0	111	~r	\ / I	\sim	1 A
1)	\/ F	- 1	\/ I	\vdash	١/١
\sim	٧ ٧	∵ ।	v i	\sim	٧.

Quantity:	100 μL
Target:	TMEM139
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TMEM139 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	TMEM139 antibody was raised using the N terminal of TMEM139 corresponding to a region
	TMEM139 antibody was raised using the N terminal of TMEM139 corresponding to a region with amino acids ITPVAYFFLTLGGFFLFAYLLVRFLEWGLRSQLQSMQTESPGPSGNARDN
Immunogen:	with amino acids ITPVAYFFLTLGGFFLFAYLLVRFLEWGLRSQLQSMQTESPGPSGNARDN
Immunogen: Specificity:	with amino acids ITPVAYFFLTLGGFFLFAYLLVRFLEWGLRSQLQSMQTESPGPSGNARDN TMEM139 antibody was raised against the N terminal of TMEM139
Immunogen: Specificity: Purification:	with amino acids ITPVAYFFLTLGGFFLFAYLLVRFLEWGLRSQLQSMQTESPGPSGNARDN TMEM139 antibody was raised against the N terminal of TMEM139
Immunogen: Specificity: Purification: Target Details	with amino acids ITPVAYFFLTLGGFFLFAYLLVRFLEWGLRSQLQSMQTESPGPSGNARDN TMEM139 antibody was raised against the N terminal of TMEM139 Affinity purified
Immunogen: Specificity: Purification: Target Details Target:	with amino acids ITPVAYFFLTLGGFFLFAYLLVRFLEWGLRSQLQSMQTESPGPSGNARDN TMEM139 antibody was raised against the N terminal of TMEM139 Affinity purified TMEM139


Application Details

Application Notes:	WB: 1 µg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	TMEM139 Blocking Peptide, catalog no. 33R-4186, is also available for use as a blocking control in assays to test for specificity of this TMEM139 antibody	
Restrictions:	For Research Use only	

Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of TMEM139 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

Images



Western Blotting

Image 1. TMEM139 antibody used at 1 ug/ml to detect target protein.