

Datasheet for ABIN633755

anti-KCNN3 antibody (C-Term)





Go to Product page

_						
	1//	Д	rv	16	٦/	٨
	W	\vdash	ΙV	Ιt	٦,	/V

Overview		
Quantity:	100 μL	
Target:	KCNN3	
Binding Specificity:	C-Term	
Reactivity:	Human, Rat, Mouse, Dog	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KCNN3 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	KCNN3 antibody was raised using the C terminal of KCNN3 corresponding to a region with	
	amino acids ITELNDRSEDLEKQIGSLESKLEHLTASFNSLPLLIADTLRQQQQQLLSA	
Specificity:	KCNN3 antibody was raised against the C terminal of KCNN3	
Purification:	Affinity purified	
Target Details		
Target:	KCNN3	
Alternative Name:	KCNN3 (KCNN3 Products)	
Background:	Action potentials in vertebrate neurons are followed by an afterhyperpolarization (AHP) that	
	may persist for several seconds and may have profound consequences for the firing pattern of	
	the neuron. Each component of the AHP is kinetically distinct and is mediated by different	

Target Details

calcium-activated potassium channels. KCNN3, a member of the KCNN family of potassium channel genes, encodes a protein that is activated before membrane hyperpolarization and is thought to regulate neuronal excitability by contributing to the slow component of synaptic AHP. The encoded protein is an integral membrane protein that forms a voltage-independent calcium-activated channel with three other calmodulin-binding subunits.

Molecular Weight:

81 kDa (MW of target protein)

Application Details

Application Notes:	WB: 2 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	KCNN3 Blocking Peptide, catalog no. 33R-4172, is also available for use as a blocking control in assays to test for specificity of this KCNN3 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of KCNN3 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.



Western Blotting

Image 1. KCNN3 antibody used at 2 ug/ml to detect target protein.