

Datasheet for ABIN634048

anti-CREG2 antibody (N-Term)





Go to Product page

_			
	IVe	rv	iew

Quantity:	100 μL	
Target:	CREG2	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CREG2 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Product Details Immunogen:	CREG2 antibody was raised using the N terminal of CREG2 corresponding to a region with	
	CREG2 antibody was raised using the N terminal of CREG2 corresponding to a region with amino acids VSSVSWAVTNEVDEELDSASTEEAMPALLEDSGSIWQQSFPASAHKEDAH	
Immunogen:	amino acids VSSVSWAVTNEVDEELDSASTEEAMPALLEDSGSIWQQSFPASAHKEDAH	
Immunogen: Specificity:	amino acids VSSVSWAVTNEVDEELDSASTEEAMPALLEDSGSIWQQSFPASAHKEDAH CREG2 antibody was raised against the N terminal of CREG2	
Immunogen: Specificity: Purification:	amino acids VSSVSWAVTNEVDEELDSASTEEAMPALLEDSGSIWQQSFPASAHKEDAH CREG2 antibody was raised against the N terminal of CREG2	
Immunogen: Specificity: Purification: Target Details	amino acids VSSVSWAVTNEVDEELDSASTEEAMPALLEDSGSIWQQSFPASAHKEDAH CREG2 antibody was raised against the N terminal of CREG2 Affinity purified	
Immunogen: Specificity: Purification: Target Details Target:	amino acids VSSVSWAVTNEVDEELDSASTEEAMPALLEDSGSIWQQSFPASAHKEDAH CREG2 antibody was raised against the N terminal of CREG2 Affinity purified CREG2	


Application Details

Application Notes:	on Notes: WB: 1 μg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	CREG2 Blocking Peptide, catalog no. 33R-9827, is also available for use as a blocking control in	
	assays to test for specificity of this CREG2 antibody	
Restrictions:	For Research Use only	

Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of CREG2 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

Images



Western Blotting

Image 1. CREG2 antibody used at 1 ug/ml to detect target protein.