

Datasheet for ABIN634061 anti-ESSPL antibody (N-Term)

1 Image



Overview

Quantity:	100 μL
Target:	ESSPL (PRSS48)
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ESSPL antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Product Details Immunogen:	ESSPL antibody was raised using the N terminal of ESSPL corresponding to a region with
	ESSPL antibody was raised using the N terminal of ESSPL corresponding to a region with amino acids MATDELATKLSRRLQMEGEGGGETPEQPGLNGAAAAAAGAPDEAAEALGS
Immunogen:	amino acids MATDELATKLSRRLQMEGEGGGETPEQPGLNGAAAAAAAAAAAAABAPDEAAEALGS
Immunogen: Specificity:	amino acids MATDELATKLSRRLQMEGEGGGETPEQPGLNGAAAAAAGAPDEAAEALGS ESSPL antibody was raised against the N terminal of ESSPL
Immunogen: Specificity: Purification:	amino acids MATDELATKLSRRLQMEGEGGGETPEQPGLNGAAAAAAGAPDEAAEALGS ESSPL antibody was raised against the N terminal of ESSPL
Immunogen: Specificity: Purification: Target Details	amino acids MATDELATKLSRRLQMEGEGGGETPEQPGLNGAAAAAAGAPDEAAEALGS ESSPL antibody was raised against the N terminal of ESSPL Affinity purified
Immunogen: Specificity: Purification: Target Details Target:	amino acids MATDELATKLSRRLQMEGEGGGETPEQPGLNGAAAAAAGAPDEAAEALGS ESSPL antibody was raised against the N terminal of ESSPL Affinity purified ESSPL (PRSS48)


Application Details

Application Notes:	WB: 1 μg/mL Optimal conditions should be determined by the investigator.
Comment:	ESSPL Blocking Peptide, catalog no. 33R-5768, is also available for use as a blocking control in assays to test for specificity of this ESSPL antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of ESSPL antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. ESSPL antibody used at 1 ug/ml to detect target protein.