

Datasheet for ABIN634139

anti-BCAT1 antibody (N-Term)





| Overview | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | BCAT1 |
| Binding Specificity: | N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BCAT1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | BCAT1 antibody was raised using the N terminal of BCAT1 corresponding to a region with |
| | amino acids MKDCSNGCSAECTGEGGSKEVVGTFKAKDLIVTPATILKEKPDPNNLVFG |
| Specificity: | BCAT1 antibody was raised against the N terminal of BCAT1 |
| Purification: | Affinity purified |
| Target Details | |
| Target: | BCAT1 |
| Alternative Name: | BCAT1 (BCAT1 Products) |
| Background: | This gene encodes the cytosolic form of the enzyme branched-chain amino acid transaminase. |
| | This enzyme catalyzes the reversible transamination of branched-chain alpha-keto acids to |
| | branched-chain L-amino acids essential for cell growth. |

Target Details

| Molecular Weight: | 43 kDa (MW of target protein) |
|---------------------|-------------------------------|
| Application Details | |

Application Notes: WB: 1 µg/mL

Optimal conditions should be determined by the investigator.

Comment: BCAT1 Blocking Peptide, catalog no. 33R-6125, is also available for use as a blocking control in


assays to test for specificity of this BCAT1 antibody

Restrictions: For Research Use only

Handling

| Format: | Lyophilized |
|------------------|--|
| Reconstitution: | Lyophilized powder. Add distilled water for a 1 mg/mL concentration of BCAT1 antibody in PBS |
| Concentration: | Lot specific |
| Buffer: | PBS |
| Handling Advice: | Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C. |

Images



Western Blotting

Image 1. BCAT1 antibody used at 1 ug/ml to detect target protein.