

Datasheet for ABIN635257

anti-DCST1 antibody (C-Term)





Go to Product page

\sim				
()	ve.	r\/	101	Λ

Quantity:	100 μL
Target:	DCST1
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DCST1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	DCST1 antibody was raised using the C terminal of DCST1 corresponding to a region with
Immunogen:	DCST1 antibody was raised using the C terminal of DCST1 corresponding to a region with amino acids SYVCRTLDCEAVYCWSCWDDMRQRCPVCTPREELSSSAFSDSNDDTAYAG
Immunogen: Specificity:	
	amino acids SYVCRTLDCEAVYCWSCWDDMRQRCPVCTPREELSSSAFSDSNDDTAYAG
Specificity:	amino acids SYVCRTLDCEAVYCWSCWDDMRQRCPVCTPREELSSSAFSDSNDDTAYAG DCST1 antibody was raised against the C terminal of DCST1
Specificity: Purification:	amino acids SYVCRTLDCEAVYCWSCWDDMRQRCPVCTPREELSSSAFSDSNDDTAYAG DCST1 antibody was raised against the C terminal of DCST1
Specificity: Purification: Target Details	amino acids SYVCRTLDCEAVYCWSCWDDMRQRCPVCTPREELSSSAFSDSNDDTAYAG DCST1 antibody was raised against the C terminal of DCST1 Affinity purified
Specificity: Purification: Target Details Target:	amino acids SYVCRTLDCEAVYCWSCWDDMRQRCPVCTPREELSSSAFSDSNDDTAYAG DCST1 antibody was raised against the C terminal of DCST1 Affinity purified DCST1


Application Details

Application Notes:	WB: 0.25 μg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	DCST1 Blocking Peptide, catalog no. 33R-8960, is also available for use as a blocking control in	
	assays to test for specificity of this DCST1 antibody	
Restrictions:	For Research Use only	

Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of DCST1 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

Images



Western Blotting

Image 1. DCST1 antibody used at 0.25 ug/ml to detect target protein.