

## Datasheet for ABIN635525 **anti-WDR33 antibody (N-Term)**



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	100 µL
Target:	WDR33
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDR33 antibody is un-conjugated
Application:	Western Blotting (WB)

#### Product Details

Immunogen:	WDR33 antibody was raised using the N terminal of WDR33 corresponding to a region with amino acids IWQRDQRDMRAIQPDAGYYNDLVPPIGMLNPMNAVTTKFVRTSTNKVKC
Specificity:	WDR33 antibody was raised against the N terminal of WDR33
Purification:	Affinity purified

#### Target Details

Target:	WDR33
Alternative Name:	WDR33 ( <a href="#">WDR33 Products</a> )
Background:	WDR33 is a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this

## Target Details

---

family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation.

Molecular Weight: 38 kDa (MW of target protein)

## Application Details

---

Application Notes: WB: 1 µg/mL  
Optimal conditions should be determined by the investigator.

Comment: WDR33 Blocking Peptide, catalog no. 33R-4214, is also available for use as a blocking control in assays to test for specificity of this WDR33 antibody

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Lyophilized powder. Add distilled water for a 1 mg/mL concentration of WDR33 antibody in PBS


Concentration: Lot specific

Buffer: PBS

Handling Advice: Avoid repeated freeze/thaw cycles.  
Dilute only prior to immediate use.

Storage: 4 °C/-20 °C

Storage Comment: Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.



### Western Blotting

**Image 1.** WDR33 antibody used at 1 ug/ml to detect target protein.