

## Datasheet for ABIN635733

# anti-NKAIN1 antibody (Middle Region)





Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:	100 μL
Target:	NKAIN1
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NKAIN1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	NKAIN1 antibody was raised using the middle region of NKAIN1 corresponding to a region with
Immunogen:	NKAIN1 antibody was raised using the middle region of NKAIN1 corresponding to a region with amino acids TPVLNSRLALEDHHVISVTGCLLDYPYIEALSSALQIFLALFGFVFACYV
Immunogen: Specificity:	
	amino acids TPVLNSRLALEDHHVISVTGCLLDYPYIEALSSALQIFLALFGFVFACYV
Specificity:	amino acids TPVLNSRLALEDHHVISVTGCLLDYPYIEALSSALQIFLALFGFVFACYV  NKAIN1 antibody was raised against the middle region of NKAIN1
Specificity: Purification:	amino acids TPVLNSRLALEDHHVISVTGCLLDYPYIEALSSALQIFLALFGFVFACYV  NKAIN1 antibody was raised against the middle region of NKAIN1
Specificity: Purification: Target Details	amino acids TPVLNSRLALEDHHVISVTGCLLDYPYIEALSSALQIFLALFGFVFACYV  NKAIN1 antibody was raised against the middle region of NKAIN1  Affinity purified
Specificity: Purification: Target Details Target:	amino acids TPVLNSRLALEDHHVISVTGCLLDYPYIEALSSALQIFLALFGFVFACYV  NKAIN1 antibody was raised against the middle region of NKAIN1  Affinity purified  NKAIN1


## **Application Details**

Application Notes:	WB: 1 µg/mL	
	Optimal conditions should be determined by the investigator.	
Comment:	NKAIN1 Blocking Peptide, catalog no. 33R-9236, is also available for use as a blocking control	
	in assays to test for specificity of this NKAIN1 antibody	
Restrictions:	For Research Use only	

## Handling

Format:	Lyophilized	
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of NKAIN1 antibody in PBS	
Concentration:	Lot specific	
Buffer:	PBS	
Handling Advice:	Avoid repeated freeze/thaw cycles.  Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.	

#### **Images**



#### **Western Blotting**

**Image 1.** NKAIN1 antibody used at 1 ug/ml to detect target protein.