

Datasheet for ABIN635880

anti-B4GALT3 antibody (N-Term)





Go to Product page

| _ | | | | |
|----|-----|-----|-----|---|
| () | ve. | rv/ | 101 | Λ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | B4GALT3 | |
| Binding Specificity: | N-Term | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This B4GALT3 antibody is un-conjugated | |
| Application: | Western Blotting (WB) | |
| Product Details | | |
| Immunogen: | B4 GALT3 antibody was raised using the N terminal of B4 ALT3 corresponding to a region with | |
| | amino acids PQGLPYCPERSPLLVGPVSVSFSPVPSLAEIVERNPRVEPGGRYRPAGCE | |
| Specificity: | B4 GALT3 antibody was raised against the N terminal of B4 ALT3 | |
| Purification: | Affinity purified | |
| Target Details | | |
| Target: | B4GALT3 | |
| Alternative Name: | B4GALT3 (B4GALT3 Products) | |
| Background: | B4GALT3 is an enzyme that may be mainly involved in the synthesis of the first N- | |
| | acetyllactosamine unit of poly-N-acetyllactosamine chains. | |
| | | |

Target Details


| Molecular Weight: | 44 kDa (MW of target protein) |
|-------------------|-------------------------------------|
| Pathways: | Glycosaminoglycan Metabolic Process |
| | |

Application Details

| Application Notes: | WB: 1 µg/mL | | |
|--------------------|--|--|--|
| | Optimal conditions should be determined by the investigator. | | |
| Comment: | B4GALT3 Blocking Peptide, catalog no. 33R-7299, is also available for use as a blocking control in assays to test for specificity of this B4GALT3 antibody | | |
| Restrictions: | For Research Use only | | |

Handling

| Format: | Lyophilized | |
|------------------|--|--|
| Reconstitution: | Lyophilized powder. Add distilled water for a 1 mg/mL concentration of B0 ALT3 antibody in PBS | |
| Concentration: | Lot specific | |
| Buffer: | PBS | |
| Handling Advice: | Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C. | |



Western Blotting

Image 1. B4GALT3 antibody used at 1 ug/ml to detect target protein.