

Datasheet for ABIN636016
anti-UXS1 antibody (Middle Region)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	UXS1
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UXS1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	UXS1 antibody was raised using the middle region of UXS1 corresponding to a region with amino acids LMLGWEPVPLEEGLNKAIHYFRKELEYQANNQYIPKPKPARIKKGRTRH
Specificity:	UXS1 antibody was raised against the middle region of UXS1
Purification:	Affinity purified

Target Details

Target:	UXS1
Alternative Name:	UXS1 (UXS1 Products)
Background:	UDP-glucuronate decarboxylase catalyzes the formation of UDP-xylose from UDP-glucuronate. UDP-xylose is then used to initiate glycosaminoglycan biosynthesis on the core protein of proteoglycans.

Target Details

Molecular Weight:	47 kDa (MW of target protein)
Pathways:	Glycosaminoglycan Metabolic Process


Application Details

Application Notes:	WB: 1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	UXS1 Blocking Peptide, catalog no. 33R-5205, is also available for use as a blocking control in assays to test for specificity of this UXS1 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Lyophilized powder. Add distilled water for a 1 mg/mL concentration of UXS1 antibody in PBS
Concentration:	Lot specific
Buffer:	PBS
Handling Advice:	Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2-8 °C for short periods. For longer periods of storage, store at -20 °C.

Images



Western Blotting

Image 1. UXS1 antibody used at 1 ug/ml to detect target protein.