

Datasheet for ABIN636622

anti-Resistin antibody



| \sim | | | | |
|--------|-------------|-----|-----|---|
| O_1 | / el | rVI | 161 | Λ |

| 0.1011 | |
|-------------------|---|
| Quantity: | 50 µg |
| Target: | Resistin (RETN) |
| Reactivity: | Human |
| Host: | Goat |
| Clonality: | Polyclonal |
| Conjugate: | This Resistin antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Blocking Antibody (Inhibition) |
| Product Details | |
| Immunogen: | Resistin antibody was raised in goat using highly pure recombinant human resistin as the |
| | immunogen. |
| Purification: | Purified |
| Target Details | |
| Target: | Resistin (RETN) |
| Alternative Name: | Resistin (RETN Products) |
| Background: | Resistin belongs to a family of tissue-specific cytokines termed FIZZ (found in inflammatory |
| | zones) and RELM. The three known members of this family, Resistin, RELM-alpha and RELM- |
| | beta share a highly conserved C-terminal domain, characterized by 10 cysteine residues with a |
| | unique spacing motif of C-X11-C-X8-C-X-C-X3-C-X10-C-X-C-X9-C-C. Resistin is an adipose- |
| | derived cytokine (adipokine) whose physiological function and molecular targets are largely |

| Target Details | |
|---------------------|---|
| | unknown. Studies have shown that Resistin suppresses insulin's ability to stimulate glucose uptake, and postulated that Resistin might be an important link between obesity and Type 2 diabetes. Other studies have indicated that Resistin expression is severely suppressed in obesity and that it may act as a feedback regulator of Adipogenesis. |
| Molecular Weight: | 20.2 kDa (predicted detection band MW) |
| Pathways: | Feeding Behaviour, Smooth Muscle Cell Migration |
| Application Details | |
| Application Notes: | Optimal conditions should be determined byt he investigator. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Concentration: | Lot specific |

| панишц | | |
|------------------|---|--|
| Format: | Lyophilized | |
| Concentration: | Lot specific | |
| Buffer: | Lyophilized from PBS, pH 7.2. | |
| Handling Advice: | Avoid repeated freeze/thaw cycles. Dilute only prior to immediate use. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | Store at -20 °C until reconstitution. Following reconstitution product may be stored at 4 °C in the short term. For long term storage aliquot and freeze at -20 °C. | |