.-online.com antibodies

Datasheet for ABIN6384870 Neurogenin 3 IQ-ELISA Kit



Overview

plasma, and serum samples.Sample Type:Cell Culture Supernatant, Plasma, SerumAnalytical Method:Semi-QuantitativeDetection Method:qPCRCharacteristics:IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a master mix are added to each well. The plate is placed into a real time PCR instrument cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.		
Reactivity:HumanMethod Type:Sandwich ELISAApplication:ELISAProduct DetailsProduct DetailsPurpose:Human Immunoquantitative (PCR-Based) Neurogenin-3 ELISA Kit for cell culture super plasma, and serum samples.Sample Type:Cell Culture Supernatant, Plasma, SerumAnalytical Method:Serni-QuantitativeDetection Method:qPCRCharacteristics:IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples be the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrument cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Quantity:	96 tests
Method Type: Sandwich ELISA Application: ELISA Product Details Purpose: Human Immunoquantitative (PCR-Based) Neurogenin-3 ELISA Kit for cell culture superplasma, and serum samples. Sample Type: Cell Culture Supernatant, Plasma, Serum Analytical Method: Semi-Quantitative Detection Method: qPCR Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumer cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Target:	Neurogenin 3 (NEUROG3)
Application: ELISA Product Details Human Immunoquantitative (PCR-Based) Neurogenin-3 ELISA Kit for cell culture superplasma, and serum samples. Sample Type: Cell Culture Supernatant, Plasma, Serum Analytical Method: Semi-Quantitative Detection Method: qPCR Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumer cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Reactivity:	Human
Product Details Purpose: Human Immunoquantitative (PCR-Based) Neurogenin-3 ELISA Kit for cell culture super plasma, and serum samples. Sample Type: Cell Culture Supernatant, Plasma, Serum Analytical Method: Semi-Quantitative Detection Method: qPCR Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standam samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a master mix are added to each well. The plate is placed into a real time PCR instrumer cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Method Type:	Sandwich ELISA
Purpose:Human Immunoquantitative (PCR-Based) Neurogenin-3 ELISA Kit for cell culture super plasma, and serum samples.Sample Type:Cell Culture Supernatant, Plasma, SerumAnalytical Method:Semi-QuantitativeDetection Method:qPCRCharacteristics:IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Application:	ELISA
plasma, and serum samples.Sample Type:Cell Culture Supernatant, Plasma, SerumAnalytical Method:Semi-QuantitativeDetection Method:qPCRCharacteristics:IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrument cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Product Details	
Sample Type: Cell Culture Supernatant, Plasma, Serum Analytical Method: Semi-Quantitative Detection Method: qPCR Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplificatio detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Purpose:	Human Immunoquantitative (PCR-Based) Neurogenin-3 ELISA Kit for cell culture supernatants,
Analytical Method: Semi-Quantitative Detection Method: qPCR Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.		plasma, and serum samples.
Detection Method: qPCR Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Sample Type:	Cell Culture Supernatant, Plasma, Serum
Characteristics: IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standard samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplificatio detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Analytical Method:	Semi-Quantitative
samples are pipetted into the wells, the target protein in the standards and samples b the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplificatio detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Detection Method:	qPCR
the immobilized antibody. The wells are washed and the detection affinity reagent is a the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplificatio detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.	Characteristics:	IQ-ELISAs employ specific capture antibodies coated on a 96-well PCR plate. Standards and
the wells where it binds to any captured antigen. The wells are washed, and primers a master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplificatio detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.		samples are pipetted into the wells, the target protein in the standards and samples binds to
master mix are added to each well. The plate is placed into a real time PCR instrumen cycling and measurement of DNA amplification. The cycle number where amplificatio detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.		the immobilized antibody. The wells are washed and the detection affinity reagent is added to
cycling and measurement of DNA amplification. The cycle number where amplification detected is proportional to the amount of affinity detection reagent that bound to capantigen in each well.		the wells where it binds to any captured antigen. The wells are washed, and primers and PCR
detected is proportional to the amount of affinity detection reagent that bound to cap antigen in each well.		master mix are added to each well. The plate is placed into a real time PCR instrument for
antigen in each well.		cycling and measurement of DNA amplification. The cycle number where amplification is
		detected is proportional to the amount of affinity detection reagent that bound to captured
Components: • Neurogenin-3 Microplate (Item A): 96 well PCR plate coated with anti-Human Neuro		antigen in each well.
	Components:	• Neurogenin-3 Microplate (Item A): 96 well PCR plate coated with anti-Human Neurogenin-3

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6384870 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

	 Wash Buffer I Concentrate (20x) (Item B): 25 ml of 20x concentrated solution Standards (Item C): 2 vials of recombinant Human Neurogenin-3 Assay Diluent A (Item D): 30 ml diluent buffer, 0.09% sodium azide as preservative. For Standard/Sample (serum/plasma) diluent Assay Diluent B (Item E): 15 ml of 5x concentrated buffer. For Standard/Sample (cell culture medium/urine) diluent Detection Affinity Reagent for Neurogenin-3 (Item F): 2 vials of a 4x concentrated solution of anti-Human Neurogenin-3 affinity reagent IQELISA Detection Reagent (Item G): 1.4ml of a 10x concentrated stock Primer Solution (Item I): 1.7 ml vial
	PCR Master Mix (Item J): 1.2 ml vial
	 PCR Preparation buffer (Item K): 1ml vial of 10x concentrated buffer
	Final Wash Buffer (Item L): 10ml vial of 10x concentrated buffer
Material not included:	Real-time PCR instrument, Bio-Rad recommended
	 Precision pipettes to deliver 2 µL to 1 mL volumes
	 Adjustable 1-25 mL pipettes for reagent preparation
	 100 mL and 1 liter graduated cylinders
	Absorbent paper
	Distilled or deionized water
	 Log-log graph paper or computer and software for data analysis
	Tubes to prepare standard or sample dilutions
	- 9

• Heating block or water bath capable of 80°C

Target Details

Target:	Neurogenin 3 (NEUROG3)
Alternative Name:	Neurogenin-3 (NEUROG3 Products)
Gene ID:	50674
UniProt:	Q9Y4Z2

Application Details

Application Notes:	The Immuno-Quantitative ELISA (IQELISA) kits are an innovative assay platform that combines
	the specificity and ease of use of an ELISA with the sensitivity of real-time PCR. This results in
	an assay that is simultaneously familiar and cutting edge and enables the use of only 1/10th
	the sample volume while also providing 10x more sensitivity than a traditional ELISA.
Comment:	The Immuno-Quantitative ELISA (IQELISA™) kits are an innovative assay platform that
	combines the specificity and ease of use of an ELISA with the sensitivity of real-time PCR. Also

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN6384870 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

	called immuno-PCR, this detection platform results in an assay that is simultaneously familiar and cutting edge. Compared to traditional ELISA, IQELISA™ enables the use of only 1/10th the sample volume while also providing 10x more sensitivity.
Sample Volume:	25 µL
Plate:	Pre-coated
Protocol:	 Prepare all reagents, samples and standards as instructed Add 25 μL standard or sample to each well. Incubate for 2.5 hours at room temperature or overnight at 4°C
	3. Add 25 µL detection affinity reagent to each well. Incubate 1 hour at room temperature
	 Add 25µL of IQELISA Detection Reagent to each well. Incubate 1 hour Add 15µL Primer solution 10µL of PCR master mix to each well
	6. Run real-time PCR
Restrictions:	For Research Use only
Handling	
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	May be stored for up to 6 months at 2° to 8°C from the date of shipment. Standard
	(recombinant protein) should be stored at -20°C or -80°C (recommended at -80°C) after
	reconstitution. Opened PCR plate or reagents may be stored for up to 1 month at 2° to 8°C.
	Note: the kit can be used within one year if the whole kit is stored at -20°C. Avoid repeated
	freeze-thaw cycles.
Expiry Date:	6 months