antibodies -online.com





MEP1A Protein (AA 22-601) (His tag)



Go to	Prod	luct	page

_							
0	V	е	r١	/	е	٧	V

Overview	
Quantity:	100 μg
Target:	MEP1A
Protein Characteristics:	AA 22-601
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This MEP1A protein is labelled with His tag.
Application:	SDS-PAGE (SDS)
Product Details	
Sequence:	MGSSHHHHHH SSGLVPRGSH MAAYKLVLIR HGESAWNLEN RFSGWYDADL SPAGHEEAKR GGQALRDAGY EFDICFTSVQ KRAIRTLWTV LDAIDQMWLP VVRTWRLNER HYGGLTGLNK
	AETAAKHGEA QVKIWRRSYD VPPPPMEPDH PFYSNISKDR RYADLTEDQL PSCESLKDTI
	ARALPFWNEE IVPQIKEGKR VLIAAHGNSL RGIVKHLEGL SEEAIMELNL PTGIPIVYEL
	DKNLKPIKPM QFLGDEETVR KAMEAVAAQG KAKK
Purity:	> 90 % by SDS - PAGE
Biological Activity Comment:	Specific activity is >300 units/mg, in which One unit will convert 1.0 umole of 3-phosphoglycerate to 2-phosphoglcerate per minute at pH 7.6 at 37C.

Target Details

Target:	MEP1A
Alternative Name:	MEP1A (MEP1A Products)
Background:	PGAM1 belongs to the phosphoglycerate mutase family. This protein is important components of glucose and 2,3-BPGA (2,3-bisphosphoglycerate) metabolism and catalyzes the reversible reaction of 3-phosphoglycerate (3-PGA) to 2-phosphoglycerate (2-PGA) in the glycolytic pathway. The PGAM is a dimeric enzyme containing, in different tissues, different proportions of a slow-migrating muscle (MM) isozyme, a fast-migrating brain (BB) isozyme, and a hybrid form (MB). Mutations in this protein cause muscle phosphoglycerate mutase efficiency, also known as glycogen storage disease X. Recombinant human PGAM protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Molecular Weight:	30.9 kDa (274aa), confirmed by MALDI-TOF
NCBI Accession:	NP_002620
UniProt:	P18669
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Liquid. In 20 mM Tris-HCl buffer(pH 8.0) containing 10 % glycerol, 1 mM DTT
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Can be stored at +4C short term (1-2 weeks). For long term storage, aliquot and store at -20C c -70C. Avoid repeated freezing and thawing cycles.