antibodies .- online.com







GST Protein (AA 1-218)



Overview

Quantity:	50 μg
Target:	GST
Protein Characteristics:	AA 1-218
Origin:	Schistosoma japonicum
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Application:	SDS-PAGE (SDS)

Product Details	
Sequence:	MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID
	GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV
	DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK
	KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPK
Purity:	> 90 % by SDS - PAGE
Endotoxin Level:	< 1.0 EU per 1ug of protein (determined by LAL method)
Biological Activity Comment:	Specific activity is > 30 units/mg, and is defined as the amount of enzyme that conjugate 1.0
	umole of 1-chloro-2,4-dinitrobenzene (CDNB) with reduced glutathione per minute at pH 6.5 at
	25C.

Target Details

Target:	GST
Alternative Name:	GST (GST Products)
Background:	GST, also known as, Glutathione S-transferase, represents a major group of detoxification enzymes. GST acts by catalyzing the reaction of glutathione with an acceptor molecule to form an S-substituted glutathione (S=sulfur). The reactions utilizing glutathione contribute the transformation of a wide range of compounds, including carcinogens, therapeutic drugs, and products of oxidative stress. As well as its enzymatic activities, GST may also bind toxins and function as transport protein. Because of this, an early term for GSTs was ligandin. GST was originally separated from Schistosomajaponicum but currently isolated from recombinant E.coli source. Recombinant Schistosoma japonicum GST was expressed in E.coli and purified by conventional chromatography techniques.
Molecular Weight:	25.4 kDa (218aa)
UniProt:	P08515

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Bioactivity Validated
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Liquid. In Phosphate Buffer Saline containing 10 % glycerol
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Can be stored at +4C short term (1-2 weeks). For long term storage, aliquot and store at -20C or -70C. Avoid repeated freezing and thawing cycles.