

Datasheet for ABIN6388273

SIRPA Protein (AA 32-373) (His tag)





Go to Product page

()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Overview	
Quantity:	100 μg
Target:	SIRPA
Protein Characteristics:	AA 32-373
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This SIRPA protein is labelled with His tag.
Product Details	

Sequence:	AA 32-373
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.

Target Details

Target:	SIRPA
Alternative Name:	SIRP alpha (SIRPA Products)
Background:	Tyrosine-protein phosphatase non-receptor type substrate 1 (SHPS1) is also known as CD172
	antigen-like family member A (CD172a), Macrophage fusion receptor, MyD-1 antigen, Signal-
	regulatory protein alpha (SIRPA or SIRP alpha) or p84, is a member of the SIRP family, and also
	belongs to the immunoglobulin superfamily. SIRP alpha is Ubiquitous and highly expressed in

brain. SIRPA / CD172a is immunoglobulin-like cell surface receptor for CD47 and acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPA / SHPS-1 supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment and may play a key role in intracellular signaling during synaptogenesis and in synaptic function By similarity. SIRPA / MyD1 involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin and mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells.

Molecular Weight:

39.8 kDa

Application Details


Restrictions:

For Research Use only

Handling


Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C

Images



ELISA

Image 1. Immobilized Mouse CD47, Mouse IgG2a Fc Tag, low endotoxin (ABIN5954914,ABIN6809967) at $2 \mu g/mL$ (100 $\mu L/well$) can bind Mouse SIRP alpha, His Tag (ABIN6386455,ABIN6388273) with a linear range of 0.02-0.625 $\mu g/mL$ (QC tested).



SDS-PAGE

 $\label{eq:mage 2.} \mbox{Mouse SIRP alpha, His Tag on under reducing (R)} \\ \mbox{condition. The gel was stained overnight with Coomassie} \\ \mbox{Blue. The purity of the protein is greater than 95 \%} \; .$