

## Datasheet for ABIN6391338

## anti-AFT1 antibody (N-Term)



Go to Product page

(	)	V		rV	ĺ	9	V	V
'	$\mathcal{I}$	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Quantity:	100 μg	
Target:	AFT1	
Binding Specificity:	N-Term	
Reactivity:	Human	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	This AFT1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	
Product Details		
Purpose:	ATF1	
Sequence:	EDSHKSTTSE TAPQ	
Isotype:	IgG	
Cross-Reactivity:	Cow, Human	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity	
	chromatography using the immunizing peptide.	
Grade:	Verified	
Target Details		
Target:	AFT1	

## **Target Details**

rarget Details				
Alternative Name:	ATF1 (AFT1 Products)			
Background:	ATF1, activating transcription factor 1, EWS-ATF1, FUS/ATF-1, TREB36, cAMP-dependent			
	transcription factor ATF-1, cyclic AMP-dependent transcription factor ATF-1			
Gene ID:	466			
NCBI Accession:	NP_005162			
Pathways:	Neurotrophin Signaling Pathway, Activation of Innate immune Response, Myometrial Relaxation			
	and Contraction, Toll-Like Receptors Cascades			
Application Details				
Application Notes:	Western Blot: Approx 37 kDa band observed in nuclear lysates of cell line A431 (calculated MW			
	of 29.2 kDa according to NP_005162.1). This molecular weight is routinely observed by other			
	sources. Recommended concentration: 0.1-0.3 µg/mL.			
	Peptide ELISA: antibody detection limit dilution 1:8000.			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	0.5 mg/mL			
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.			
Preservative:	Sodium azide			
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which			
	should be handled by trained staff only.			
Handling Advice:	Minimize freezing and thawing.			
Storage:	-20 °C			
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated			
	at 4°C for a few weeks and still remain viable.			