# antibodies -online.com





## anti-NAP1L1 antibody (C-Term)

2 Images



Go to Product page

$\sim$				
	11/6	٦r١	/10	۱۸.

Quantity:	400 μL	
Target:	NAP1L1	
Binding Specificity:	AA 352-381, C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NAP1L1 antibody is un-conjugated	
Application:	Western Blotting (WB), Flow Cytometry (FACS)	
Product Details		
Immunogen:	This NAP1L1 antibody is generated from rabbits immunized with a KLH conjugated synthetic	
	peptide between 352-381 amino acids from the C-terminal region of human NAP1L1.	
Clone:	RB8412	
Isotype:	Ig Fraction	
Predicted Reactivity:	B, M, Rat	
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.	
Target Details		
Target:	NAP1L1	

#### **Target Details**


Alternative Name:	NAP1L1 (NAP1L1 Products)	
Background:	NAP1L1 is a member of the nucleosome assembly protein (NAP) family. This protein participates in DNA replication and may play a role in modulating chromatin formation and contribute to the regulation of cell proliferation.	
Molecular Weight:	45374	
Gene ID:	4673	
NCBI Accession:	NP_004528, NP_631946	
UniProt:	P55209	

#### **Application Details**

Application Notes:	WB: 1:1000. FC: 1:10~50
Restrictions:	For Research Use only


#### Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



### Flow Cytometry

**Image 1.** NL1 Antibody (C-term) 920a flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goatanti-rabbit secondary antibodies were used for the analysis.



#### **Western Blotting**

**Image 2.** Western blot analysis of NL1 (arrow) using rabbit polyclonal NL1 Antibody 920a. 293 cell lysates ( $2 \mu g$ /lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the NL1 gene.