

Datasheet for ABIN651208
anti-CCDC92 antibody (AA 177-205)[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	CCDC92
Binding Specificity:	AA 177-205
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC92 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This CCD92 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 177-205 amino acids from the Central region of human CCD92.
Clone:	RB26053
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	CCDC92
Alternative Name:	CCD92 (CCDC92 Products)
Molecular Weight:	36961

Target Details

Gene ID:	80212
NCBI Accession:	NP_079416
UniProt:	Q53HC0


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. CCD92 Antibody (Center) (ABIN651208 and ABIN2840130) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CCD92 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of CCD92 Antibody (Center) (ABIN651208 and ABIN2840130) in mouse testis tissue lysates (35 µg/lane). CCD92 (arrow) was detected using the purified Pab.