

Datasheet for ABIN651234
anti-IN80D antibody (N-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	400 µL
Target:	IN80D (INO80D)
Binding Specificity:	AA 63-91, N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IN80D antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This IN80D antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 63-91 amino acids from the N-terminal region of human IN80D.
Clone:	RB26035
Isotype:	Ig Fraction
Predicted Reactivity:	X
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	IN80D (INO80D)
Alternative Name:	IN80D (INO80D Products)

Target Details

Molecular Weight:	113202
Gene ID:	54891
NCBI Accession:	NP_060229
UniProt:	Q53TQ3


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. IN80D Antibody (N-term) (ABIN651234 and ABIN2840146) IHC analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the IN80D Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of IN80D Antibody (N-term) (ABIN651234 and ABIN2840146) in mouse liver tissue lysates (35 µg/lane).IN80D (arrow) was detected using the purified Pab.