



[Go to Product page](#)

Datasheet for ABIN651274

anti-TPRG1L antibody (N-Term)

2 Images

Overview

Quantity:	400 µL
Target:	TPRG1L
Binding Specificity:	AA 35-63, N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TPRG1L antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This TPRGL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 35-63 amino acids from the N-terminal region of human TPRGL.
Clone:	RB26064
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	TPRG1L
Alternative Name:	TPRGL (TPRG1L Products)
Molecular Weight:	30212

Target Details

Gene ID: 127262

NCBI Accession: [NP_877429](#)

UniProt: [Q5T0D9](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:50~100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. TPRGL Antibody (N-term) (ABIN651274 and ABIN2840164) IHC analysis in formalin fixed and paraffin embedded human cervix carcinoma tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TPRGL Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. Western blot analysis of TPRGL Antibody (N-term) (ABIN651274 and ABIN2840164) in mouse bladder tissue lysates (35 µg/lane).TPRGL (arrow) was detected using the purified Pab.