antibodies -online.com





anti-TTPAL antibody (C-Term)

3 Images



Go to Product page

_					
U	V	er	VI	е	W

Quantity:	400 μL
Target:	TTPAL
Binding Specificity:	AA 204-233, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TTPAL antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Product Details Immunogen:	This TTPAL antibody is generated from rabbits immunized with a KLH conjugated synthetic
	This TTPAL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 204-233 amino acids from the C-terminal region of human TTPAL.
Immunogen:	peptide between 204-233 amino acids from the C-terminal region of human TTPAL.
Immunogen: Clone:	peptide between 204-233 amino acids from the C-terminal region of human TTPAL. RB26315
Immunogen: Clone: Isotype:	peptide between 204-233 amino acids from the C-terminal region of human TTPAL. RB26315 Ig Fraction
Immunogen: Clone: Isotype: Purification:	peptide between 204-233 amino acids from the C-terminal region of human TTPAL. RB26315 Ig Fraction

Target Details

Background:	TTPAL may act as a protein that binds a hydrophobic ligand.
Gene ID:	79183
NCBI Accession:	NP_001034288, NP_077307
UniProt:	Q9BTX7


Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

Handling


Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.	
Expiry Date:	6 months	

Images




Flow Cytometry

Image 1. TTL Antibody (C-term) (ABIN651357 and ABIN2840199) flow cytometric analysis of MCF-7 cells (right histogram) comred to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western Blotting

Image 2. Western blot analysis of TTL Antibody (C-term) (ABIN651357 and ABIN2840199) in MCF-7cell line lysates $(35 \, \mu g/lane)$.TTL (arrow) was detected using the purified b.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. TTL Antibody (C-term) (ABIN651357 and ABIN2840199) IHC analysis in formalin fixed and raffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TTL Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.