

Datasheet for ABIN651560  
**anti-GGNBP2 antibody (C-Term)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	400 µL
Target:	GGNBP2
Binding Specificity:	AA 647-676, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GGNBP2 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	This GGNBP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 647-676 amino acids from the C-terminal region of human GGNBP2.
Clone:	RB26137
Isotype:	Ig Fraction
Predicted Reactivity:	C, M
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	GGNBP2
---------	--------

## Target Details



Alternative Name:	GGNBP2 ( <a href="#">GGNBP2 Products</a> )
Molecular Weight:	79086
Gene ID:	79893
NCBI Accession:	<a href="#">NP_079111</a>
UniProt:	<a href="#">Q9H3C7</a>

## Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.
Expiry Date:	6 months



### Flow Cytometry

**Image 1.** GGNBP2 Antibody (C-term) (ABIN651560 and ABIN2840301) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### Western Blotting

**Image 2.** Anti-GGNBP2 Antibody (C-term) at 1:1000 dilution + Jurkat whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.

### Immunohistochemistry (Paraffin-embedded Sections)

**Image 3.** GGNBP2 Antibody (C-term) (ABIN651560 and ABIN2840301) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GGNBP2 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.