# antibodies -online.com





# anti-ATP11C antibody (AA 589-616)

4 Images



Publication



Go to Product page

Overview	
Quantity:	400 μL
Target:	ATP11C
Binding Specificity:	AA 589-616
Reactivity:	Human, Mouse, Hamster
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATP11C antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded
	Sections) (IHC (p))
Product Details	
Product Details Immunogen:	This ATP11C antibody is generated from rabbits immunized with a KLH conjugated synthetic
	This ATP11C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 589-616 amino acids from the Central region of human ATP11C.
Immunogen:	peptide between 589-616 amino acids from the Central region of human ATP11C.
Immunogen: Clone:	peptide between 589-616 amino acids from the Central region of human ATP11C.  RB22082
Immunogen:  Clone: Isotype:	peptide between 589-616 amino acids from the Central region of human ATP11C.  RB22082  Ig Fraction
Immunogen:  Clone:  Isotype:  Purification:	peptide between 589-616 amino acids from the Central region of human ATP11C.  RB22082  Ig Fraction

## **Target Details**

rarget Details	
Background:	The function of ATP11C remains unknown.
Molecular Weight:	129477
Gene ID:	286410
NCBI Accession:	NP_001010986, NP_775965
UniProt:	Q8NB49
Application Details	
Application Notes:	WB: 1:1000. WB: 1:1000. IHC-P: 1:50~100. FC: 1:10~50
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Maintain refrigerated at 2-8 °C for up to 6 months. For long term storage store at -20 °C in small aliquots to prevent freeze-thaw cycles.


### **Publications**

6 months

Expiry Date:


Product cited in:

Hu, Zhou, Zhao, Wu: "Integrin  $\alpha$ 6/Akt/Erk signaling is essential for human breast cancer resistance to radiotherapy." in: **Scientific reports**, Vol. 6, pp. 33376, (2018) (PubMed).




#### **Flow Cytometry**

**Image 1.** ATP11C Antibody (Center) (ABIN651638 and ABIN2840341) flow cytometric analysis of 293 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



#### **Western Blotting**

**Image 2.** ATP11C Antibody (Center) (ABIN651638 and ABIN2840341) western blot analysis in mouse lung tissue lysates ( $35 \,\mu g/lane$ ). This demonstrates the ATP11C antibody detected ATP11C protein (arrow).



#### **Western Blotting**

**Image 3.** ATP11C Antibody (Center) (ABIN651638 and ABIN2840341) western blot analysis in CHO cell line lysates (35  $\mu$ g/lane).This demonstrates the ATP11C antibody detected the ATP11C protein (arrow).

Please check the product details page for more images. Overall 4 images are available for ABIN651638.